Further Pure Mathematics FP1

Candidates sitting Further Pure Mathematics FP1 may also require those formulae listed under Pure Mathematics P1 and P2.

Summations

$$\sum_{r=1}^{n} r^2 = \frac{1}{6} n(n + 1)(2n + 1)$$

$$\sum_{r=1}^{n} r^3 = \frac{1}{4} n^2(n + 1)^2$$

Numerical solution of equations

The Newton-Raphson iteration for solving $f(x) = 0$: $x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)}$

Conics

	Parabola	**Rectangular Hyperbola**
Standard Form	$y^2 = 4ax$	$xy = c^2$
Parametric Form	$(at^2, 2at)$	$\left(ct, \frac{c}{t}\right)$
Foci	$(a, 0)$	Not required
Directrices	$x = -a$	Not required

Matrix transformations

Anticlockwise rotation through θ about O: $\begin{pmatrix} \cos\theta & -\sin\theta \\ \sin\theta & \cos\theta \end{pmatrix}$

Reflection in the line $y = (\tan\theta)x$: $\begin{pmatrix} \cos 2\theta & \sin 2\theta \\ \sin 2\theta & -\cos 2\theta \end{pmatrix}$

Pearson Edexcel International Advanced Subsidiary/Advanced Level in Mathematics, Further Mathematics and Pure Mathematics – Mathematical Formulae and Statistical Tables – Issue 2 – January 2021 © Pearson Education Limited 2021

Pure Mathematics P1

Mensuration

Surface area of sphere $= 4\pi r^2$

Area of curved surface of cone $= \pi r \times$ slant height

Cosine rule

$a^2 = b^2 + c^2 - 2bc\cos A$

Pure Mathematics P2

Arithmetic series

$u_n = a + (n-1)d$

$S_n = \frac{1}{2}n(a+l) = \frac{1}{2}n[2a + (n-1)d]$

Geometric series

$u_n = ar^{n-1}$

$S_n = \frac{a(1-r^n)}{1-r}$

$S_\infty = \frac{a}{1-r}$ for $|r| < 1$

Logarithms and exponentials

$\log_a x = \frac{\log_b x}{\log_b a}$

Binomial series

$$(a+b)^n = a^n + \binom{n}{1}a^{n-1}b + \binom{n}{2}a^{n-2}b^2 + \ldots + \binom{n}{r}a^{n-r}b^r + \ldots + b^n \quad (n \in \mathbb{N})$$

where $\binom{n}{r} = {}^nC_r = \frac{n!}{r!(n-r)!}$

$$(1+x)^n = 1 + nx + \frac{n(n-1)}{1\times 2}x^2 + \ldots + \frac{n(n-1)\ldots(n-r+1)}{1\times 2\times \ldots \times r}x^r + \ldots \quad (|x| < 1, n \in \mathbb{R})$$

Numerical integration

The trapezium rule: $\int_a^b y \, dx \approx \frac{1}{2}h\{(y_0 + y_n) + 2(y_1 + y_2 + \ldots + y_{n-1})\}$, where $h = \frac{b-a}{n}$